


Assessment and Management of Climate Change in the Chesapeake TMDL

2022 National Training Workshop: Water Quality Data, Assessment, and Plans

June 2, 2022

Lew Linker (EPA-CBP) and the CBPO Modeling Team

llinker@chesapeakebay.net



Chesapeake Bay Program
Science, Restoration, Partnership




The CBP Climate Change Assessment


- The CBP has developed the tools to quantify the effects of three decades of climate change (1995 to 2025) on Chesapeake water quality standards through changes in watershed flows and loads, storm intensity, estuarine temperatures, sea level rise, and ecosystem influences including loss of tidal wetland attenuation with sea level rise.
- Using the climate change assessment tools, the CBP decided in December 2020 to implement management practices to meet new nutrient and sediment targets that address current climate change in the Chesapeake watershed by 2025.
- Current efforts are to develop new assessment tools for the airshed, watershed and tidal estuary to estimate 2035 climate change risk to Chesapeake TMDL water quality standards.




Overview of Bay Designated Uses



A. Cross Section of Chesapeake Bay or Tidal Tributary




B. Oblique View of the "Chesapeake Bay" and its Tidal Tributaries





Components of Climate Change – Effect on Tidal Dissolved Oxygen






Elements of Chesapeake Water Quality Climate Risk Assessment

Climate Change Processes and Dependencies

Model
Data Set
Endpoint
Project/Decision

- Complete
- In Process
- Not included But important
- Not included minor

Initials indicate the responsible person
Numbers indicate the section of the documentation



Climate

Watershed

Estuary

Management

Approaches, Methods, and Findings from the Watershed



Chesapeake Bay Program
Science, Restoration, Partnership



Analysis of Climate Change in the Chesapeake Watershed

For the analysis of climate change in the Chesapeake watershed, the primary components considered are precipitation volume, precipitation intensity, temperature, and evapotranspiration with an additional consideration to CO₂ concentrations.

Overall, increased precipitation volumes and intensity are estimated to increase nutrient and sediment loads from the watershed in 2025, 2035, 2045, and 2055 compared to 1995.



However, increased future temperatures substantially ameliorates the effect of estimated increased precipitation volume in the watershed through evapotranspiration.



Chesapeake Bay Program
Science, Restoration, Partnership

For the 2025 Climate Change Estimate:

The trends in annual precipitation on a county level were developed through the application of PRISM data and analysis provided and recommended by Jason Lynch, EPA, and Karen Rice, USGS. The annual PRISM dataset for the years 1927 to 2014 (88 years) were used in for the regression trend analysis. For the analysis PRISM data were first spatially aggregated for each Phase 6 land segments. The Phase 6 land segments typically represent a county. For each land segment a simple linear trend was fitted to the annual rainfall dataset.




Annual rainfall volumes for the 88-year period linear regression lines are shown in red for the two land segments (counties) – (a) Centre County in Pennsylvania and (b) District of Columbia. The values for the slope of the regression lines, and the corresponding 30-year projections in the rainfall volume (1995 to 2025) are also shown.

Source: Section 12 of Phase 6 Documentation



Assessment of Influence of 2025 Climate Change in the Watershed




Projections of rainfall increase using trend in 88-years of annual PRISM^[1] data

Change in Rainfall Volume 2021-2030 vs. 1991-2000

2025 Rainfall Projection (percent change)

- 0.7 - 1.0
- 1.1 - 1.5
- 1.6 - 2.0
- 2.1 - 2.5
- 2.6 - 3.0
- 3.1 - 3.5
- 3.6 - 4.0
- 4.1 - 4.5
- 4.6 - 5.0
- 5.1 - 5.5
- 5.6 - 6.0




Major Basins	PRISM Trend
Youghiogheny River	2.1%
Patuxent River Basin	3.3%
Western Shore	4.1%
Rappahannock River Basin	3.2%
York River Basin	2.6%
Eastern Shore	2.5%
James River Basin	2.2%
Potomac River Basin	2.8%
Susquehanna River Basin	3.7%
Chesapeake Bay Watershed	3.1%

[1] Parameter-elevation Relationships on Independent Slopes Model



Trends in Observed Rainfall Intensity



Observed changes in rainfall intensity in the Chesapeake region over the last century. The equal allocation distribution (blue) is contrasted with the distribution obtained based on observed changes (red).




1940-2014 streamflow trends based on observations

Chesapeake Bay Program
Science, Restoration, Partnership

The study analyzed USGS GAGES-II data for a subset of Hydro-Climatic Data Network 2009 (HCDN-2009).


Annual Average Streamflow in the United States, 1940-2014



More than 50% decrease 20% to 50% decrease 20% decrease to 20% increase 20% to 50% increase More than 50% increase

Data source: USGS (U.S. Geological Survey), 2016. Analysis of data from the National Water Information System. Accessed May 2016.

For more information, visit U.S. EPA's "Climate Change Indicators in the United States" at www.epa.gov/climate-indicators.



Karen C. Rice, Douglas L. Moyer, and Aaron L. Mills, 2017. Riverine discharges to Chesapeake Bay: Analysis of long-term (1927 - 2014) records and implications for future flows in the Chesapeake Bay basin *JEM* 204 (2017) 246-254

USGS station ID	Precipitation		Discharge	
	Slope	p-value	Slope	p-value
04252500	0.0007	0.0011	0.0021	<0.0001
01512500	0.0008	0.0007	0.0016	0.0028
01503000	0.0007	0.0022	0.0013	0.0181
01531000	0.0006	0.0219	0.0018	0.0030
01531500	0.0007	0.0044	0.0016	0.0029
01532000	0.0006	0.0374	0.0015	0.0330
01534000	0.0005	0.0497	0.0015	0.0120
01550000	0.0005	0.0493	0.0019	0.0015
01543000	0.0004	0.1000	0.0018	0.0058
01545500	0.0004	0.0953	0.0017	0.0026
01536500	0.0006	0.0078	0.0016	0.0027
01551500	0.0005	0.0612	0.0017	0.0017
01439500	0.0005	0.0972	0.0007	0.1661
01541500	0.0003	0.2357	0.0017	0.0017
01540500	0.0006	0.0111	0.0016	0.0023
01541000	0.0004	0.0985	0.0016	0.0021
01567000	0.0004	0.1577	0.0011	0.0250
01570500	0.0005	0.0260	0.0013	0.0088

North-South Split

01562000	0.0004	0.1693	0.0007	0.2082
01638500	0.0004	0.1150	0.0008	0.1026
01608500	0.0004	0.1725	0.0010	0.0833
01636500	0.0005	0.1245	0.0008	0.0624
01606500	0.0003	0.1958	0.0009	0.1108
01668000	0.0006	0.0794	0.0004	0.4727
02035000	0.0003	0.2653	-0.0001	0.8243
02019500	0.0002	0.4333	0.0003	0.4836
03488000	0.0003	0.2480	0.0006	0.2841

U.S. Environmental Protection Agency. 2016. Climate change indicators in the United States, 2016. Fourth edition. EPA 430-R-16-004. www.epa.gov/climate-indicators.

Annual average percent change were calculated using Sen slope (Helsel and Hirsch, 2002).

Lins, H.F. 2012. USGS Hydro-Climatic Data Network 2009 (HCDN-2009). U.S. Geological Survey Fact Sheet 2012-3047. <https://pubs.usgs.gov/fs/2012/3047>.

Helsel, D.R., and R.M. Hirsch. 2002. Statistical methods in water resources. Techniques of water resources investigations, Book 4. Chap. A3. U.S. Geological Survey. <https://pubs.usgs.gov/twri/twri4a3>.

An ensemble of GCM projections from BCSD CMIP5^[1] was used to estimate 1995-2025 temperature change.

?

Data Unavailable

GCM Used

Selection updated

Updated Ensemble members		
ACCESS1-0	FGOALS-g2	IPSL-CM5A-LR
BCC-CSM1-1	FIO-ESM	IPSL-CM5A-MR
BCC-CSM1-1-M	GFDL-CM3	IPSL-CM5B-LR
BNU-ESM	GFDL-ESM2G	MIROC-ESM
CanESM2	GFDL-ESM2M	MIROC-ESM-CHEM
CCSM4	GISS-E2-H-CC	MIROC5
CESM1-BGC	GISS-E2-R	MPI-ESM-LR
CESM1-CAM5	GISS-E2-R-CC	MPI-ESM-MR
CMCC-CM	HadGEM2-AO	MRI-CGCM3
CNRM-CM5	HadGEM2-CC	NorESM1-M
CSIRO-MK3-6-0	HadGEM2-ES	31 member ensemble
EC-EARTH	INMCM4	

[1] BCSD – Bias Correction Spatial Disaggregation;
[1] CMIP5 – Coupled Model Intercomparison Project 5

Reclamation, 2013. 'Downscaled CMIP3 and CMIP5 Climate and Hydrology Projections: Release of Downscaled CMIP5 Climate Projections, Comparison with preceding Information, and Summary of User Needs', prepared by the U.S. Department of the Interior, Bureau of Reclamation, Technical Services Center, Denver, Colorado. 47pp.

Source: Kyle Hinson, VIMS




Chesapeake Bay Program
Science, Restoration,
Partnership


Chesapeake Bay Watershed Annual Change in Temperature

Degrees Celsius

2025 - RCP 4.5




1.1°C Increase
in Annual
Temperature




Degrees Celsius

2050 - RCP 4.5









1.94°C Increase
in Annual
Temperature






Temperature trends for the six CBP states

Annual temperature for 1895 to 2015 are shown.





Approx. increases over the last 30 years based on the trend line are shown.



NOAA National Climatic Data Center
<https://www.ncdc.noaa.gov/temp-and-precip/state-temps/>



Estimated potential evapotranspiration




(a) Relative change in estimated change in potential evapotranspiration due to change in temperature is shown from different methods. It shows temperature alone can introduce considerable differences in estimation of potential evapotranspiration with the selection of method. (b) Estimate of percent changes in potential evapotranspiration




Estimates of Climate Only and Climate and Land Use


Marginal Differences in Freshwater Delivery




Marginal Differences in Sediment Delivery



Marginal Differences in Nitrogen Delivery



Marginal Differences in Phosphorus Delivery



Grey bar = climate only Black bar = Climate and Land Use



Elements of 2025 Climate Change (1995-2025)


**Air-temperature
increase: 1.06 °C**



↑
**Sea Level
Rise:
0.22m**

**Open boundary:
Temperature: +0.95 °C;
Salinity: +0.18 psu
(Thomas et al., 2017)**

- Flow
→ 2.4% Increase
- Nitrogen Load
→ 2.6% Increase
- Phosphorus Load
→ 4.5% Increase
- Sediment Load
→ 3.8% Increase



Phase 6 Watershed Model

**Model: CH3D-ICM
400m-1km Resolution**




Elements of 2035 Climate Change (1995-2035)

**Air-temperature
increase: 1.39 °C**

**Percent Change in Rainfall (2035 vs. 1995)
Hybrid of Extrapolation and GCMs**

- 2.4% - 3.2%
- 3.3% - 3.8%
- 3.9% - 4.2%
- 4.3% - 4.8%
- 4.9% - 5.5%
- 5.6% - 6.3%

**+4.21 %
Increase
in rainfall**



Phase 6 Watershed Model

Flow

→
3.7% Increase

Nitrogen Load

→
4.7% Increase

Phosphorus Load

→
9.9% Increase

Sediment Load

→
8.5% Increase



**Model: CH3D-ICM
400m-1km Resolution**

**Sea Level
Rise:
0.31m**



**Open boundary:
Temperature: +1.32 °C;
Salinity: +0.25 psu
(Thomas et al., 2017)**

Approaches, Methods, and Findings from the Tidal Bay




Chesapeake Bay Program
Science, Restoration, Partnership



Analysis of Climate Change in the Tidal Bay

Estimates of the influence of sea level rise, increased temperature of tidal waters, and tidal wetland loss were incorporated into the Water Quality and Sediment Transport Model (WQSTM) of the tidal Bay (Cercio and Noel 2017). Guidance for increasing levels of regional sea level rise based upon global tide gauge rates and regional land subsidence rates came from the Climate Resiliency Workgroup CRWG). Specifically, the CRWG recommended that sea level rise projections for 2025 be based on long term observations at Sewells Point, VA (0.22 m).







Bottom DO Change: 1995 to 2025

Keeping all other factors constant, sea level rise and increased watershed flow reduce hypoxia in the Bay, but the predominant influence are the negative impacts of increased water column temperature.


Sea Level Rise




Watershed Flow




Increased Temp.



All Factors



Summer (Jun.-Sep.) Hypoxia Volume (<1 mg/l) 1991-2000 in the Whole Bay Under 2025 WIP3 Condition



Estimated 2025 Climate Change Influence

**Air-temperature
increase: 1.06 °C**

Sea Level Rise: 0.22m

Flow

+2.4% est. 2025

TN

+2.6% est. 2025

TP


+4.5% est. 2025

Sediment

+3.8 est. 2025




Tidal wetland change



Open boundary delta T: + 0.95 °C; delta S: + 0.18 psu
(Thomas et al., 2017)



Chesapeake Bay Tidal Wetlands




Source: Carl Cerco, U.S. CoE ERDC

- The extent from National Wetlands Inventory is determined largely from vegetation perceived via aerial photography.
- 190,000 hectares of estuarine (green) and tidal fresh (red) wetlands.
- A tidal wetlands module is now fully operational in the WQSTM. The module incorporates functions of sediment and particulate nutrient removal and burial, denitrification, and respiration. The loss of wetland function due to sea level rise and inundation will be accounted for explicitly.



Influence of Estimated 2025 (0.3 m) and 2050 (0.5m) Sea Level Rise on Tidal Wetland Attenuation



There is little change in estimated total tidal wetland area for 2025 (0.17 - 0.3 m) and 2050 (0.5 m) which equates to negligible changes in tidal wetland attenuation.

Long range (2100) conditions estimate tidal wetland changes to be on the order of a 40% loss in the Chesapeake which could reduce tidal wetland attenuation on the order of about 10 million pounds nitrogen and 0.6 million pounds phosphorus.




Δ Achievement of Deep Channel DO Water Quality Standard

Chesapeake Bay Program
Science, Restoration, Partnership



Achievement of **Deep Channel DO** water quality standard (1mg/l instantaneous minimum) expressed as **an incremental increase** over the PSC agreed to 2025 planning targets

CB Segment	State	2025 Climate	2035 Climate	2045 Climate	2055 Climate
		2025 Land Use	2025 Land Use	2025 Land Use	2025 Land Use
		204TN	208TN	212TN	220TN
		14.0TP	14.6TP	15.4TP	16.7TP
		1993-1995	1993-1995	1993-1995	1993-1995
		DO Deep Channel	DO Deep Channel	DO Deep Channel	DO Deep Channel
CB3MH	MD	0.00%	0.00%	0.00%	0.00%
CB4MH	MD	1.47%	3.15%	4.62%	7.31%
CB5MH	MD	0.00%	0.00%	0.00%	0.00%
CB5MH	VA	0.00%	0.00%	0.00%	0.00%
POTMH	MD	0.00%	0.00%	0.00%	0.00%
RPPMH	VA	0.00%	0.00%	0.00%	0.00%
ELIPH	VA	0.00%	0.00%	0.00%	0.00%
CHSMH	MD	0.01%	0.92%	1.08%	2.34%





Climate Target Loads in Perspective




Overall, the CBP found that a target load of 5 million pounds nitrogen and 0.6 million pounds phosphorus will be sufficient to offset 30 years of climate change in the Chesapeake Bay.

Model load reduction estimates from CAST-2019 (current version of the CBP watershed model)

Summer Hypoxia Volume Averaged Over Ten Years for All Tidal Waters

June - Sept. with Critical DO Values of 1 mg/l and 3 mg/l





Conclusions:

Climate change is a multigenerational challenge for the CBP and is a force multiplier for headwinds to the Chesapeake restoration.

However, the CBP is working on management practices that are effective counters to climate change such as:

- The design and accelerated adoption of stormwater management practices appropriately designed for increased rainfall volumes and intensities that are expected in the future for all counties in the Chesapeake watershed.
- Examination of the top tier agriculture and urban BMPs that are most vulnerable to future climate risk, with an emphasis on practices that could be adapted to become more resilient to future climate conditions of increased rainfall intensities and volumes.
- A quantification of the co-benefits of BMPs that mitigate future climate risk as they relate to the protection of local infrastructure, public health and safety, green infrastructure, urban floodplain management, riparian buffers, tidal and non-tidal wetlands, cold water fisheries, and other management actions.

The climate change risk to the Chesapeake TMDL can be effectively managed and the CBP is actively addressing the challenge.